

# Species richness, geographical affinities and activity patterns of mammals in premontane Andean forests of the Magdalena River basin of Colombia

Diego A. Torres<sup>1</sup>, Abel Eduardo Rojas<sup>1</sup>

<sup>1</sup> Facultad de Ciencias Exactas y Naturales, Universidad de Caldas, Calle 65 # 26-10, Manizales, Colombia

Corresponding author: Diego A. Torres ([dtorresarboleda@gmail.com](mailto:dtorresarboleda@gmail.com))

Academic editor: A.M. Leal-Zanchet | Received 31 July 2020 | Accepted 25 January 2021 | Published 23 February 2021

Citation: Torres DA, Rojas AE (2021) Species richness, geographical affinities and activity patterns of mammals in premontane Andean forests of the Magdalena River basin of Colombia. *Neotropical Biology and Conservation* 16(1): 145–166. <https://doi.org/10.3897/neotropical.16.e57109>

## Abstract

More than half of the population of Colombia is settled in the Magdalena River basin, resulting in high deforestation rates due to productive activities and urbanisation. Within this scenario of forest loss and ecosystem degradation, it is imperative to record and monitor the biodiversity in order to decrease and mitigate the negative consequences of human activities on species and ecosystems. For six years, we assessed the mammal species richness, abundance and activity patterns in premontane forests of the Magdalena River basin in the Department of Caldas, Colombia. We also presented additional information on the geographical affinities of this fauna. We recorded 101 species, seven of them endemic to Colombia, with Chiroptera being the richest order, followed by Rodentia. Most of the species are common and not listed in threatened categories and only four are vulnerable and two endangered, according to the Red List of the IUCN and the Ministerio de Medio Ambiente y Desarrollo Sostenible of Colombia. The mammalian fauna of the study area is similar to that of other lowland localities in the Neotropics and different to the fauna in highland localities, including the nearby ones. Specifically, this fauna was most similar to that in lowland Tolima and the Caribbean Region of Colombia, Venezuela and Costa Rica; however, when we accounted only for bat fauna, it was more similar to the fauna in Caribbean and Pacific Regions of Colombia. To secure the long-term persistence of these species, we recommend maintenance of the current corridors, such as riparian forests and living fences and an increase in the forested area.

## Keywords

biodiversity, biogeography, Caldas, checklist, inventory, monitoring

## Introduction

The Magdalena River basin is located in northern South America in Colombia (Morrone 2014). This river has its origin in the Andes mountains (Colombian Massif) and flows northwards between the Central and Eastern Cordilleras throughout the Magdalena River Valley, emptying finally into the Caribbean Sea (Hermelin 2016). Biogeographically, this basin is classified as the Magdalena Province and belongs to the Pacific domain, sharing a taxa composition with nearby Provinces, such as Guajira, Chocó-Darién and Cauca (Hernández-Camacho et al. 1992; Kattan et al. 2004; Morrone 2014).

This Basin is of great economic relevance for the country because a large part of the Colombian population and their productive activities are settled there (IDEAM 2001). Such a level of economic development has produced a continuous pressure on natural resources, resulting in high rates of deforestation mainly at low to mid elevations (Armenteras et al. 2013). It is estimated that around 70% of Andean and 30% of lowland forests in Colombia have been lost (Etter et al. 2006a, b), especially in the Magdalena River basin (Etter and van Wyngaarden 2000). The current landscapes in this basin are very heterogeneous, where forests are represented mainly by fragments immersed in an agricultural matrix with different levels of connection through riparian forests and living fences (Etter et al. 2008). The remaining extensive areas of forests are mainly associated with the Andean highlands, natural parks and private civil reserves, as well as hydroelectric dams (Armenteras et al. 2003).

In this context of forest loss and fragmentation, it is important to monitor the biodiversity in order to prevent, mitigate, compensate or correct the negative effects of socio-economic activities on ecosystems (Schmeller et al. 2017). An important tool for this purpose is species inventories which result in checklists, natural history field data and spatiotemporal trends in species richness and abundance (Christoffersen 2010, Lees et al. 2014). The data obtained from the previous activities allow the construction and improvement of species distributional hypotheses (Elith et al. 2006) and to the identification of evidence-based important areas for conservation (Niemelä 2000).

In the Magdalena River and its tributaries, mammal inventories can be tracked back as far as the 19<sup>th</sup> century (Mantilla-Meluk et al. 2014) and, during the last decades, checklists have been frequently published (e.g. Moreno-Bejarano and Álvarez-León 2003; Castaño and Corrales 2010; Garcés Retrepo et al. 2016; Solari et al. 2020). However, most published inventories and monitoring are based on one year or less sampling efforts and the local mammalian richness is generally underestimated. Here, we provide a checklist for mammals from premontane forests on the eastern slopes of the Central Cordillera in the Department of Caldas, Colombia, as a result of field data collection distributed over six years (2014–2019). We also provide some insights into the geographical affinities of this fauna and data on their activity patterns.

## Methods

### Study area

The forests, included in this study, are distributed on the eastern slopes of the Central Cordillera in the Department of Caldas, Colombia. We surveyed forests in the Municipalities of Victoria, Norcasia and Samaná in an elevation range between 300 and 1000 metres. This heterogeneous landscape is composed of crops, pastures and natural vegetation, ranging from stubble to riparian forests and mature secondary forests. All sampling sites were located in the basin of the rivers Manso, Miel and Guarinó. These last two are tributaries of the Magdalena River. Sampling sites were under the influence of the area around the Miel I hydroelectric dam. The average temperature was 23 °C with a maximum of 33 °C with warmer conditions at lower elevations. Annual average precipitation varies from year to year between 3000 to 5000 mm and distributed in an annual bimodal pattern with December to February and June to August as the dry periods (IDEAM 2001).

### Mammal sampling

We accumulated 647 sampling days during the six years (2014–2019). The samplings were distributed during both rainy and dry seasons. To capture bats, we used mist-nets installed in the understorey, across streams and at forest edges. Mist-nets remained open after sunset until 22:00 h. Manual captures were opportunistic, mainly associated with species roosting under small bridges, in hollow trunks or in the foliage. For mist-nets, a total of 34000 metres of net-nights was accumulated. For small and medium non-volant mammals sampling, we used live capture traps (Sherman and Tomahawk), located on the ground and up to two metres above the ground on branches. Traps were baited with a mixture of banana and oat, flavoured with vanilla essence or sardine and corn with bacon butter. Each trap was checked daily in the morning and the bait replaced with fresh bait. The traps were installed in linear transects by stations 10–15 m apart, for a total of 20–25 stations according to the available area at each sampling site. Each station contained a trap on the ground and a trap in the branches. In total, we accumulated 33455 trap-nights. Mammals were also sampled using trap-cameras (Bushnell) located along trails and streams where the passage of mammals was highly probable. Cameras were set 300 m apart at a minimum and we accumulated a total of 3435 hours-cam. Direct observations were also included.

### Mammalian taxonomy and conservation status

For bat taxonomy, we followed Simmons and Cirranello (2020). For taxonomic updates in *Chiroderma* and *Tonatia*, we followed Lim et al. (2020) and Basantes et al. (2020). We treated *Glossophaga soricina* on the western side of the Andes as a differ-

ent species (here *Glossophaga* sp.), as suggested by Hoffmann et al (2019). We also treated the *Sturnira* in the study area as an undetermined species because some authors classify it as *S. parvidens* or *S. giannae* (García-Herrera et al. 2019; Esquivel et al. 2020); however, the distributions of these species are expected to be restricted to Central America and the Amazon basin, respectively (Velazco and Patterson 2019; Hernández-Canchola and León-Paniagua 2020). Thus, the populations in our study area cannot yet be confidently assigned to any species.

The taxonomy of non-volant mammals followed the Mammal Taxonomy Database of the American Society of Mammalogists (Burgin et al. 2018). For squirrels, we followed the taxonomic arrangement proposed in Fiedler et al. (2020) and for *Marmosa*, we followed Voss et al. (2020). Some specimens were collected, prepared as skull and skin and deposited in the Museo de Historia Natural of the Universidad de Caldas. All procedures followed the guidelines of the American Society of Mammalogists for the use of wild mammals in research (Sikes and Gannon 2011). The global status of conservation and population trends followed the Red List of the IUCN ([www.iucnredlist.org](http://www.iucnredlist.org)). The national conservation status followed the Resolución 1912 of 2017 of the Ministerio de Ambiente y Desarrollo Sostenible.

## Data analysis

To establish geographical similarities of the mammalian fauna in the study area with other areas of the Neotropics, we constructed a matrix of presence/absence of 548 species of mammals from data available from checklists for localities in Central America and northern South America (Suppl. material 1). We excluded the genera *Mazama* and *Sylvilagus* because it was not clear to what species to assign the species reported in the references. We gathered data for the following areas: (1) the Central Cordillera of Colombia on the western slopes in Risaralda (Castaño et al. 2018), Valle del Cauca (Rojas-Díaz et al. 2012), Cauca (Ramírez-Chaves and Pérez 2010) and the eastern slopes in Tolima (García-Herrera et al. 2019); (2) the Western Cordillera of Colombia on both slopes at the Departments of Valle del Cauca and Cauca and the Pacific lowlands (Ramírez-Chaves and Pérez 2010; Rojas-Díaz et al. 2012); (3) the eastern slopes of the Colombian Massif in Cauca (Ramírez-Chaves and Pérez 2010); (4) the Caribbean Region of Colombia in Córdoba (Race-ro-Casarrubia et al. 2015); (5) the Orinoquía of Colombia in Arauca (Mosquera-Guerra et al. 2019); (6) the Guiana Shield in Colombia (Trujillo et al. 2018) and (7) French Guiana (Lim 2012); (8) the Amazonas in Venezuela (Lim 2012); (9) the Sierra de Aroa in Yarucay State in northern Venezuela (García et al. 2016); (10) Central America in Costa Rica (Rodríguez-Herrera et al. 2014); and (11) La Rioja in Argentina (Fariñas-Torres et al. 2018), this last locality being chosen to be used as outgroup.

For the Colombian Cordilleras, we grouped separately the mammalian fauna below 1000 m from that recorded over 2000 m of elevation to analyse them as different localities because highland mammalian fauna in the Andes tend to be different

from that in the lowlands (Mena et al. 2011; Velazco and Patterson 2013). Localities were clustered, based on the Jaccard Similarity Index using the algorithm Paired Group (UPGMA) in the software PAST (Hammer et al. 2001).

We assessed inventory completeness as  $RO/RE \times 100$ , where RO was the observed species richness and RE was the species richness estimated by the index Chao 1, calculated with the software ESTIMATES, based on a matrix of presence or absence of species and randomised 100 times (Colwell and Elsensohn 2014) using days as sampling units. Finally, as we did not intend to assess habitat use, but only activity patterns, we used raw total abundances obtained in camera traps, constructing a frequency distribution graph for each of the 24 hours of the day for species with 20 or more records. We presented a checklist for the area, including two additional species (*Mustela frenata* and *Centronycteris centralis*) recorded for the area by Castaño and Corrales (2007 and 2010), but not recorded in this study.

## Results


### Mammal richness

We gathered 9848 records of mammals (recaptures not included) representing 101 species from nine orders and 26 families (Table 1; Figs 1–4). Chiroptera was the richest order with 53 species, followed by Rodentia with 19 species. Two species are listed globally as Endangered and one as Vulnerable and four species are listed nationally as Vulnerable. Seven species are endemic to Colombia, most of them rodents (four species). The completeness of the inventory was 91.5% (Fig. 5).

### Geographical affinities


The mammal fauna in the study area was most similar to the fauna on the eastern slopes of the Central Cordillera in Tolima below 1000 m (Fig. 6A). Concurrently, the fauna of these two areas was similar to the one of the Caribbean Region in Córdoba. Together, the fauna of these three areas grouped with that of the Sierra de Aroa in Venezuela and Costa Rica and formed a group with the fauna of the Venezuelan Amazon and Guiana Shield in Colombia and French Guiana. Altogether, this group was similar to another group containing the fauna of Central and Western cordilleras of Colombia (Fig. 6A).

When only bat richness was considered, the pattern of similarity amongst localities changes (Fig. 6B). Bat fauna in the study area was also more similar to the fauna of Tolima below 1000 m and both fauna were similar to those of the Caribbean in Córdoba. However, instead of grouping with Central America or the Sierra de Aroa, the bat fauna of these three areas was more similar to that in the Pacific area of Valle del Cauca and Cauca (Colombia). All fauna making up this group were similar to other groups formed by the bat fauna of Venezuelan Amazon and Guiana Shield (Fig. 6B).




**Figure 1.** Some bat species captured between 2014 and 2019 in premontane forests of the Magdalena River basin in eastern Caldas, Colombia. *Artibeus amplus* (A), *A. phaeotis* (B), *A. lituratus* (C), *A. ravus* (D), *A. planirostris* (E), *Carollia brevicauda* (F), *C. castanea* (G), *C. perspicillata* (H), *Phyllostomus hastatus* (I), *P. discolor* (J), *Phylloderma stenops* (K), *Lonchorhina aurita* (L), *Lophostoma brasiliense* (M), *L. silvicolum* (N), *Tonatia bakeri* (O), *Trinycteris nicefori* (P), *Chiroderma gorgasi* (Q), *Mesophylla macconnelli* (R), *Vampyressa thuyone* (S), *Platyrhinus helleri* (T), *Uroderma convexum* (U), *Sturnira* sp. (V), *Vampyrum spectrum* (W) and *Lamproncycteris brachyotis* (X).





**Figure 2.** Some bat species captured between 2014 and 2019 in premontane forests of the Magdalena River basin in eastern Caldas, Colombia. *Lonchophylla robusta* (A), *Glossophaga* sp. (B), *Anoura caudifer* (C), *A. geoffroyi* (D), *Lichonycteris* aff. *obscura* (E), *Micronycteris microtis* (F), *M. schmidtorum* (G), *Desmodus rotundus* (H), *Cormura brevirostris* (I), *Rhynchonycteris naso* (J), *Saccopteryx bilineata* (K), *Noctilio leporinus* (L), *Eptesicus chiriquinus* (M), *E. brasiliensis* (N), *Myotis riparius* (O), *Rhogeessa io* (P), *Cynomops greenhalli* (Q), *Eumops hansae* (R) and *Molossus bondae* (S).




**Figure 3.** Some marsupials and rodents captured between 2014 and 2019 in premontane forests of the Magdalena River basin in eastern Caldas, Colombia. *Caluromys lanatus* (A), *Didelphis marsupialis* (B), *Metachirus myosuros* (C), *Marmosa isthmica* (D), *M. phaea* (E), *Syntheosciurus granatensis* (F), *Tylomys mirae* (G), *Transandinomys talamancae* (H), *Zygodontomys* aff. *brunneus* (I), *Rhipidomys caucensis* (J), *Melanomys caliginosus* (K), *Handleyomys alfaroi* (L), *Proechimys chrysaеolus* (M) and *Nectomys grandis* (N).


### Activity patterns

Mammals showed three types of activity pattern (Fig. 7): (1) diurnal, including species as *Dasyprocta punctata* and *Eira barbara*; (2) cathemeral (active both at day and night), such as *Procyon cancrivorus*; and (3) nocturnal, like the rest of the analysed species. However, some species showed activities that fell out of their regular activity pattern; for example, *E. barbara* showed activity in the first hours after dawn and *Tamandua mexicana* and *Proechimys chrysaеolus* showed activity during the morning and noon. All the 10 analysed species showed some crepuscular activity.






**Figure 4.** Some mammals registered between 2014 and 2019 in premontane forests of the Magdalena River basin in eastern Caldas, Colombia. *Leopardus pardalis* (A), *Procyon cancrivorus* (B), *Cercopithecus* (C), *Galictis vittata* (D), *Pecari tajacu* (E), *Cabassous centralis* (F), *Saguinus leucopus* (G) and *Aotus griseimembra* (H).




**Figure 5.** Curves of estimated (Chao 1) and observed species richness between 2014 and 2019 in premontane forests of the Magdalena River basin in eastern Caldas, Colombia.

## A. All mammals




## B. Bats



**Figure 6.** Dendrograms showing the similarity in species composition (Jaccard Index) of mammals (A) and bats (B) amongst northern South America and Central America localities. Red star indicates the study area. Central, West and East refer to the three Andes Cordilleras in Colombia; W (west) and E (east) refers to the slopes of the Andes; SA (South America); CA (Caribbean); Central-East refers to the Colombian Massif.

## Discussion

Premontane forests, located on the eastern slopes of the Central Andes below 1000 m in the Department of Caldas, sustain at least 101 species of mammals, mostly common species that are not listed as threatened. This number represented 19% of the mammalian richness in Colombia (528 species; Sociedad Colombiana



**Figure 7.** Activity patterns of some mammals between 2014 and 2019 in premontane forests of the Magdalena River basin in eastern Caldas, Colombia. Dark grey areas indicate hours of darkness, while light grey indicates twilight. Records represent the observations in trap-cameras during the six years of monitoring.

**Table 1.** Checklist of mammals in premontane forests of the Magdalena River basin in eastern Caldas, Colombia.

ORDER/FAMILY	Conservation status			Elevation	Records	Method	Municipality
	National	Global	Population trend	m			
DIDELPHIMORPHIA							
Didelphidae							
<i>Caluromys lanatus</i>		LC	Decreasing	494–850	10	Cam, Trap, Obs	Nor, Sam
<i>Chironectes minimus</i>		LC	Decreasing	501–816	16	Obs, Trap	Nor, Sam, Vic
<i>Didelphis marsupialis</i>		LC	Stable	303–880	245	Cam, MC, Trap, Obs	Nor, Sam, Vic
<i>Marmosa phaea</i>		LC	Stable	334–848	31	MC, Trap, Obs	Nor, Sam, Vic
<i>Marmosa isthmica</i>		Not evaluated	Unknown	324–867	84	Cam, MC, Trap	Nor, Sam, Vic
<i>Marmosa robinsoni</i>		LC	Stable	687–825	2	Trap	Sam
<i>*Marmosops chucha</i>		Not evaluated	Unknown	450–808	17	MC, Trap	Nor, Sam, Vic
<i>Metachirus myosuros</i>		LC	Stable	461–860	98	Cam, MC, Trap, Obs	Nor, Sam, Vic
<i>Monodelphis adusta</i>		LC	Stable	542–795	3	Trap	Nor, Vic
<i>Philander melanurus</i>		LC	Stable	781	1	Cam	Vic
CINGULATA							
Dasyopodidae							
<i>Cabassous centralis</i>		DD	Unknown	532–860	12	Cam, MC	Nor, Sam
<i>Dasyus novemcinctus</i>		LC	Stable	394–850	27	Cam, Obs	Nor, Sam, Vic
PILOSA							
Megalonychidae							
<i>Choloepus hoffmanni</i>		LC	Unknown	528–817	4	Obs	Nor, Sam
Myrmecophagidae							
<i>Tamandua mexicana</i>		LC	Unknown	487–860	35	Cam, Obs	Nor, Sam, Vic
CHIROPTERA							
Emballonuridae							
<i>Centronycteris centralis</i>		LC	Unknown	420	–	Castaño and Corrales (2007)	Nor
<i>Cormura brevirostris</i>		LC	Unknown	518–591	5	Obs, MisN	Nor, Sam
<i>Peropteryx macrotis</i>		LC	Stable	378	1	MisN	Vic
<i>Rhynchonycteris naso</i>		LC	Unknown	675	1	MisN	Sam
<i>Saccopteryx bilineata</i>		LC	Unknown	376–820	9	Obs, MisN	Nor, Sam, Vic
<i>Saccopteryx leptura</i>		LC	Unknown	468–686	6	MisN	Nor, Sam
Molossidae							
<i>Cynomops greenhallii</i>		LC	Unknown	675	4	MisN	Sam
<i>Eumops hansae</i>		LC	Unknown	675	2	MisN	Sam
<i>Molossus bondae</i>		LC	Stable	675	6	MisN	Sam
Noctilionidae							
<i>Noctilio albiventris</i>		LC	Stable	661	1	MisN	Sam
<i>Noctilio leporinus</i>		LC	Unknown	503–675	2	MisN	Sam, Vic
Phyllostomidae							
<i>Anoura caudifer</i>		LC	Unknown	379–822	7	MisN	Nor, Sam, Vic
<i>Anoura geoffroyi</i>		LC	Stable	819	3	MisN	Nor
<i>Artibeus amplus</i>		LC	Unknown	372–873	59	MisN	Nor, Sam, Vic
<i>Artibeus bogotensis</i>		LC	Stable	1000	5	MisN	Sam
<i>Artibeus lituratus</i>		LC	Stable	408–857	239	MisN	Nor, Sam, Vic
<i>Artibeus phaeotis</i>		LC	Unknown	353–873	207	MisN	Nor, Sam, Vic
<i>Artibeus planirostris</i>		LC	Stable	372–857	272	MisN	Nor, Sam, Vic
<i>Artibeus rufus</i>		LC	Stable	408–873	225	MisN	Nor, Sam, Vic
<i>Carollia brevicauda</i>		LC	Stable	353–887	1007	MisN	Nor, Sam, Vic
<i>Carollia castanea</i>		LC	Stable	353–887	973	MisN	Nor, Sam, Vic
<i>Carollia perspicillata</i>		LC	Stable	353–887	2693	MisN	Nor, Sam, Vic
<i>Chiroderma salvini</i>		LC	Stable	666	1	MisN	Sam
<i>Chiroderma gorgasi</i>		Not evaluated	Unknown	584–799	2	MisN	Nor, Sam
<i>Choeroniscus</i> aff. <i>minor</i>		LC	Unknown	794	1	MisN	Nor
<i>Desmodus rotundus</i>		LC	Stable	353–840	131	MisN	Nor, Sam, Vic
<i>Glossophaga</i> sp.		LC	Stable	408–812	17	MisN	Nor, Vic
<i>Lampronnycteris brachyotis</i>		LC	Stable	372–666	12	MisN	Sam, Vic
<i>Lichonycteris</i> aff. <i>obscura</i>		LC	Unknown	654–778	13	MisN	Sam, Vic



ORDER/FAMILY	Conservation status			Elevation m	Records	Method	Municipality
	National	Global	Population trend				
<i>Lonchophylla robusta</i>		LC	Unknown	478–819	8	MisN	Nor, Sam
<i>Lonchorhina aurita</i>		LC	Stable	372–694	100	MisN	Nor, Sam, Vic
<i>Lophostoma brasiliense</i>		LC	Stable	493–849	39	MC, MisN	Nor, Sam, Vic
<i>Lophostoma silvicolium</i>		LC	Unknown	511–668	2	MisN	Sam
<i>Mesophylla macconnelli</i>		LC	Unknown	493–845	66	MisN	Nor, Sam, Vic
<i>Micronycteris hirsuta</i>		LC	Unknown	830	1	MisN	Sam
<i>Micronycteris megalotis</i>		LC	Unknown	372	1	MisN	Vic
<i>Micronycteris microtis</i>		LC	Stable	518	2	MisN	Nor
<i>Micronycteris minuta</i>		LC	Unknown	373–763	2	MisN	Nor, Vic
<i>Micronycteris schmidtorum</i>		LC	Stable	468–853	10	MisN	Nor, Sam
<i>Phyllostoma stenops</i>		LC	Stable	373–745	12	MisN	Nor, Sam, Vic
<i>Phyllostoma discolor</i>		LC	Stable	372–830	34	MisN	Nor, Sam, Vic
<i>Phyllostoma hastatus</i>		LC	Stable	372–830	19	MisN	Nor, Sam, Vic
<i>Platyrrhinus helleri</i>		LC	Stable	372–887	279	MisN	Nor, Sam, Vic
<i>Sturmira ludovici</i>		LC	Unknown	651–661	2	MisN	Sam
<i>Sturmira sp.</i>				372–887	771	MisN	Nor, Sam, Vic
<i>Tonatia bakeri</i>		Not evaluated	Unknown	468–644	18	MisN	Nor, Sam, Vic
<i>Trinycteris nicefori</i>		LC	Unknown	353–585	3	MisN	Sam, Vic
<i>Uroderma convexum</i>		Not evaluated	Unknown	376–873	37	MisN	Nor, Sam, Vic
<i>Vampyressa thuyone</i>		LC	Unknown	478–853	126	MisN	Nor, Sam, Vic
<i>Vampyrus spectrum</i>		NT	Decreasing	505	2	MisN	Nor
<b>Thyropteridae</b>							
<i>Thyroptera tricolor</i>		LC	Unknown	502–742	10	MC, MisN	Sam, Vic
<b>Vespertilionidae</b>							
<i>Eptesicus brasiliensis</i>		LC	Unknown	814–826	5	MisN	Nor, Sam
<i>Eptesicus chiroquimus</i>		LC	Unknown	805–819	2	MisN	Nor, Sam
<i>Myotis riparius</i>		LC	Stable	372–887	99	MisN	Nor, Sam, Vic
<i>Rhogeessa io</i>		LC	Unknown	567–846	2	MisN	Nor, Sam
<b>CARNIVORA</b>							
<b>Canidae</b>							
<i>Cerdocyon thous</i>		LC	Stable	461–871	27	Cam, Obs	Nor, Sam, Vic
<b>Felidae</b>							
<i>Leopardus pardalis</i>		LC	Decreasing	475–864	22	Cam, Obs	Nor, Sam
<b>Mustelidae</b>							
<i>Eira barbara</i>		LC	Decreasing	389–868	61	Cam, Obs	Nor, Sam, Vic
<i>Galictis vittata</i>		LC	Stable	748–814	4	Cam	Nor, Sam
<i>Lontra longicaudis</i>	Vu	NT	Decreasing	475–585	4	Obs, Tra	Nor, Sam
<i>Mustela frenata</i>		LC	Stable	–	–	Castañó and Corrales (2010)	Nor
<b>Procyonidae</b>							
<i>Nasua nasua</i>		LC	Decreasing	666	1	Cam	Sam
<i>Potos flavus</i>		LC	Decreasing	295–843	19	Obs	Nor, Sam, Vic
<i>Procyon cancrivorus</i>		LC	Decreasing	486–887	61	Cam, Obs	Nor, Sam, Vic
<b>ARTIODACTYLA</b>							
<b>Tayassuidae</b>							
<i>Pecari tajacu</i>		LC	Stable	499–692	42	Cam	Sam, Vic
<b>Primates</b>							
<b>Atelidae</b>							
<i>Alouatta seniculus</i>		Not evaluated	Unknown	460	1	Obs	Vic
<b>Cebidae</b>							
<i>Aotus griseimembra</i>	Vu	Vu	Decreasing	416–853	132	Obs	Nor, Sam, Vic
<i>Saguinus leucopus</i>	Vu	EN	Decreasing	372–887	424	Cam, Obs	Nor, Sam, Vic
<i>*Cebus versicolor</i>		EN	Decreasing	1000	1	Obs	Sam
<b>RODENTIA</b>							
<b>Cricetidae</b>							
<i>Handleyomys alfaroi</i>		LC	Stable	320–828	20	Trap	Nor, Sam
<i>Ichthyomys hydrobates</i>		NT	Decreasing	797–800	2	Obs, Trap	Nor, Sam
<i>Melanomys caliginosus</i>		LC	Stable	333–804	65	Trap	Nor, Sam, Vic
<i>Neacomys tenuipes</i>		LC	Stable	491–891	64	Cam, MC, Trap, Obs	Nor, Sam, Vic

ORDER/FAMILY	Conservation status			Elevation m	Records	Method	Municipality
	National	Global	Population trend				
<i>*Nectomys grandis</i>		DD	Unknown	658–826	17	Trap	Nor, Sam
<i>*Rhipidomys caucensis</i>		DD	Unknown	509–781	10	Trap	Nor, Sam
<i>Rhipidomys latimanus</i>				334–801	3	Trap	Nor, Vic
<i>Sigmodon hirsutus</i>		LC	Increasing	475–808	3	Trap	Nor, Vic
<i>Transandinomys talamancae</i>		LC	Stable	790–825	8	Trap	Sam
<i>Tylomys mirae</i>		LC	Unknown	314–891	124	Cam, MC, Trap, Obs	Nor, Sam, Vic
<i>*Zygodontomys aff. brunneus</i>		LC	Stable	475–521	15	Trap	Nor
<b>Cuniculidae</b>							
<i>Cuniculus paca</i>		LC	Stable	488–838	92	Cam, Obs	Nor, Sam, Vic
<b>Dasyproctidae</b>							
<i>Dasyprocta punctata</i>		LC	Stable	390–860	268	Cam, Obs	Nor, Sam, Vic
<b>Dinomysidae</b>							
<i>Dinomys branickii</i>	Vu	LC	Unknown	849	1	Obs	Vic
<b>Echimyidae</b>							
<i>*Proechimys chrysaeolus</i>		DD	Unknown	303–829	161	Cam, Obs, Trap	Nor, Sam, Vic
<b>Erethizontidae</b>							
<i>Coendou quichua</i>		DD	Decreasing	510	1	Obs	Nor
<b>Heteromyidae</b>							
<i>Heteromys australis</i>		LC	Stable	730–840	5	Trap	Sam
<b>Sciuridae</b>							
<i>Syntheosciurus granatensis</i>		LC	Stable	460–887	36	Cam, Obs	Nor, Sam, Vic
<i>Leptosciurus pucheranii</i>		DD	Unknown	468–888	13	Cam, MC, Obs	Nor, Sam
<b>LAGOMORPHA</b>							
<b>Leporidae</b>							
<i>Sylvilagus sp.</i>		–	–	497	1	Cam	Vic

National (resolución 1912 de 2017) and global (Red List of IUCN) conservation status: **DD** (data deficient), **LC** (least concern), **NT** (near threatened), **VU** (vulnerable), **EN** (endangered). Methods: **Cam** (camera trap), **MC** (manual capture), **MisN** (mist net), **Obs** (direct observation), **Trap** (Sherman and Tomahawk traps). Municipalities: **Nor** (Norcasia), **Sam** (Samaná), **Vic** (Victoria).

de Mastozoología et al. 2020) and it is expected to increase with additional sampling efforts. Bats as the richest order and rodents as the second follow the same pattern found at the national scale (Solari et al. 2013).

We expected to find, according to the estimator Chao 1, at least nine species more if the sampling effort were increased. Checklists of mammals in localities in the Magdalena River basin report species that were not found in the study area, but their presence is highly probable, especially bats such as *Myotis caucensis*, *M. albes-cens*, *Pteronotus parnellii*, *Trachops cirrhosus*, some molossid species and nectarivorous bats, such as *Hsunitycteris thomasi* and *Lionycteris spurrelli* (García-Herera et al. 2019; Solari et al. 2020). Some carnivorous mammals, such as *Herpailurus yagouaroundi* or *Bassaricyon neblina*, are also expected (Sánchez-Giraldo and Daza 2017; Gerstner et al. 2018). Besides increasing the sampling effort, it is important to focus it on habitats where rare species concentrate their activities. This was the case of the bat *Eumops hansae*, which was captured over the Manso River in a net set across the river, a method that was rarely used in this or other studies; the specimen captured by this method was the first reported for the country (Torres and Rojas 2020).

The mammalian fauna found in the study area is composed mostly of common species with wide geographic ranges; however, the presence of seven Colombian endemic small mammals with restricted geographic ranges associated with premon-tane and cloud forests in the Magdalena River basin emphasises the conservation

value of premontane forests in the east of the Department of Caldas; indeed, this area had already been identified as a regional centre of mammalian endemisms (Castaño 2012). Since non-volant small mammals have low dispersal ability, local deforestation associated with agricultural activities can cause local population extinctions of these endemic species and restrict them to isolated forested areas (Castro and Fernandez 2004, Paise et al. 2020). This is especially true for the marsupial *Marmosops chucha*, the rodent *Rhipidomys caucensis* and the primate *Saguinus leucopus* that carry out their activities above the ground in the lower, middle and upper forest strata (pers. obs., Poveda and Sánchez-Palomino 2004). To ensure the survival in the long term for these endemic species, it is imperative to increase the connectivity of forest fragments and maintain the already existing corridors such as riparian forests and living fences (Zimbres et al. 2017).

Most species in the study area are not listed in lower threatened categories (LC or NT); however, 10 species showed a globally-decreasing population trend. This pattern is part of a world phenomenon known as defaunation, which means the progressive loss of animal populations caused mainly by habitat loss (Dirzo et al. 2014). Interestingly, eight of the eleven species with decreasing populations, according to the Red List, feed on other animals (i.e. animalivorous), despite belonging to different taxonomic groups, such as bats (*Vampyrus spectrum*), carnivores (*Leopardus pardalis* or *Lontra longicaudis*), marsupials (*Chironectes minimus*) and rodents (*Ichthyomys hydrobates*). Animalivory is a functional trait of species that has been associated with increased vulnerability to habitat loss and fragmentation because of their lower population densities and slow life histories (Cardillo et al. 2004; Minin et al. 2016).

Since ambient temperature is an important determinant of the distribution of many animals (Bozinovic et al. 2011) and the inter-Andean valley, through which the Magdalena River flows, opens in the north, the faunistic similarity between the study area and the warm Caribbean Region in Colombia, Costa Rica and Venezuela was not surprising. This inter-Andean valley could be defined as a “warm corridor” for mammalian species adapted to these warm environmental conditions. The warm lowlands similarity was also supported by the fact that mammalian fauna in the Central Cordillera below 1000 m (both the study area and Tolima) and the Caribbean localities are more related to the warm lowlands of the Amazon and Guiana Shield, than they are to closer regions like West and Central Cordilleras above 2000 m, which have lower temperatures. Our analysis restricted to bats also points to this relationship and adds to the group of “lowland fauna”, the bat fauna of the Pacific Region below 1000 metres.

Similar relationships between lowland areas have also been described for birds (Kattan et al. 2004; Cadena et al. 2016). For these flying vertebrates, it has been hypothesised that relationships amongst lowland areas in northern South America were mediated by dispersal events (Cadena et al. 2016). One hypothesis is that this dispersion occurred throughout the north in the Caribbean lowlands (Haffer 1967a, b). Another hypothesis proposes that this occurred through passes in the Andes, such

as the Táchira depression in northern East Cordillera or Andalucía pass and Suaza-Pescado valleys at the south of this same Cordillera (Cadena et al. 2016). Possible examples of this historical dispersion in mammals may be the bats *Artibeus planirostris* and *Sphaeronycteris toxophyllum* which are distributed in the lowlands east of the Andes, but also extend their distribution into the Magdalena River basin and Caribbean Region (Solari et al. 2013; García-Herrera et al. 2018). At least in the case of *A. planirostris*, molecular evidence suggests dispersion from the Guiana Shield and Venezuela into the Lesser Antilles and northern South America (Larsen et al. 2007).

Most analysed mammal species were active during the night, beginning at dusk and finishing at sunrise. This is the typical mammalian pattern; and, as it is currently understood, this was the ancestral behaviour of the placental mammalian ancestor (Gerkema et al. 2013); thus, activity during the day, as shown by some species, may be considered a derived behavioural trait. In general, the activity patterns of many Neotropical mammals of medium and large size are well-known (e.g. Blake et al. 2012; Ramírez-Mejía and Sánchez 2016; Huck et al. 2017); however, the small mammals activity patterns are under-studied (Ferreira and Vieira 2016). Thus, the data, shown in this study on the activity patterns of the endemic spiny rat *Proechimys chrysaеolus*, are an important contribution to this topic. The activity pattern of *P. chrysaеolus* was similar to the patterns found in other species of *Proechimys* in the Brazilian Amazon (Pratas-Santiago et al. 2016).

## Conclusion

Premontane forests of the Magdalena River basin in eastern Caldas harbour a rich mammalian fauna, consisting mostly of common species of lowland origins. However, these forests are of high conservation value because they host at least seven endemic species and five Endangered or Vulnerable species. To secure the long-term persistence of these species, we recommend maintaining the current corridors, such as riparian forests and living fences and increasing the forested area.

## Acknowledgements

ISAGEN and the Universidad de Caldas financed this study (agreement 33/45). We thank Beatriz Edilma Toro, John Harold Castaño and Thomas Defler for valuable comments to this manuscript and Mary Luz Bedoya for administrative and logistical support. We are grateful to editors and two anonymous reviewers who greatly improved this manuscript.

## References

- Armenteras D, Cabrera E, Rodríguez N, Retana J (2013) National and regional determinants of tropical deforestation in Colombia. *Regional Environmental Change* 13: 1181–1193. <https://doi.org/10.1007/s10113-013-0433-7>



- Armenteras D, Gast F, Villareal H (2003) Andean forest fragmentation and the representativeness of protected natural areas in the eastern Andes, Colombia. *Biological Conservation* 113: 245–256. [https://doi.org/10.1016/S0006-3207\(02\)00359-2](https://doi.org/10.1016/S0006-3207(02)00359-2)
- Basantes M, Tinoco N, Velazco PM, Hofmann MJ, Rodríguez-Posada ME, Camacho MA (2020) Systematics and taxonomy of *Tonatia saurophila* Koopman & Williams, 1951 (Chiroptera, Phyllostomidae). *ZooKeys* 915: 59–86. <https://doi.org/10.3897/zookeys.915.46995>
- Blake JG, Mosquera D, Loiselle BA, Swing K, Guerra J, Romo D (2012) Temporal activity patterns of terrestrial mammals in lowland rainforest of Eastern Ecuador. *Ecotropica* 18: 137–146.
- Bozinovic F, Calosi P, Spicer JJ (2011) Physiological correlates of geographic range in animals. *Annual Review of Ecology Evolution and Systematics* 42: 155–179. <https://doi.org/10.1146/annurev-ecolsys-102710-145055>
- Burgin CJ, Colella JP, Kahn PL, Upham NS (2018) How many species of mammals are there? *Journal of Mammalogy* 99: 1–14. <https://doi.org/10.1093/jmammal/gyz052>
- Cadena CD, Pedraza CA, Brumfield RT (2016) Climate, habitat associations and the potential distributions of Neotropical birds: Implications for diversification across the Andes. *Revista de la Academia Colombiana de Ciencias Exactas, Físicas y Naturales* 40: 275–287. <https://doi.org/10.18257/raccefyn.280>
- Cardillo M, Purvis A, Sechrest W, Gittleman JL, Bielby J, Mace GM (2004) Human population density and extinction risk in the world's carnivores. *PLoS Biology* 2: 909–914. <https://doi.org/10.1371/journal.pbio.0020197>
- Castaño JH, Corrales JD (2010) Mamíferos de la Cuenca del río la Miel (Caldas): diversidad y uso cultural. *Boletín Científico, Centro de Museos, Museo de Historia Natural* 14: 56–75. [http://www.scielo.org.co/scielo.php?pid=S0123-30682010000100004&script=sci\\_abstract&tlng=es](http://www.scielo.org.co/scielo.php?pid=S0123-30682010000100004&script=sci_abstract&tlng=es)
- Castaño JH, Torres DA, Rojas-Díaz V, Saavedra-Rodríguez CA, Pérez-Torres J (2018) Mamíferos del departamento de Risaralda, Colombia. *Biota Colombiana* 18: 240–255. <https://doi.org/10.21068/c2017.v18n02a16>
- Christoffersen ML (2010) Continental biodiversity of South American oligochaetes: The importance of inventories. *Acta Zoológica Mexicana* 26: 35–46. <https://doi.org/10.21829/azm.2010.262876>
- Colwell RK, Elsensohn JE (2014) EstimateS turns 20: Statistical estimation of species richness and shared species from samples, with non-parametric extrapolation. *Ecography* 37: 609–613. <https://doi.org/10.1111/ecog.00814>
- Dirzo R, Young H, Galetti M, Ceballos G, Isaac NJB, Collen B (2014) Defaunation in the anthropocene. *Science* 345: 401–406. <https://doi.org/10.1126/science.1251817>
- Elith J, Graham CH, Anderson RP, Dudík M, Ferrier S, Guisan A, Hijmans RJ, Huettmann F, Leathwick JR, Lehmann A, Li J, Lohmann LG, Loiselle BA, Manion G, Moritz C, Nakamura M, Nakazawa Y, Overton JMM, Peterson AT, Phillips SJ, Richardson K, Scachetti-Pereira R, Schapire RE, Soberón J, Williams S, Wisz MS, Zimmermann NE (2006) Novel methods improve prediction of species' distributions from occurrence data. *Ecography* 29: 129–151. <https://doi.org/10.1111/j.2006.0906-7590.04596.x>

- Esquivel DA, Aya-Cuero CA, Peña S, Velásquez-Guarín D, Ramírez-Chaves HE (2020) Murciélagos (Chiroptera) del departamento del Tolima, Colombia: diversidad en un bosque húmedo tropical y actualización de la lista de especies del departamento. *Boletín científico Centro de museos Museo de historia natural* 24: 71–99.
- Etter A, McAlpine C, Phinn S, Pullar D, Possingham H (2006a) Unplanned land clearing of Colombian rainforests: Spreading like disease? *Landscape and Urban Planning* 77: 240–254. <https://doi.org/10.1016/j.landurbplan.2005.03.002>
- Etter A, McAlpine C, Pullar D, Possingham H (2006b) Modelling the conversion of Colombian lowland ecosystems since 1940: Drivers, patterns and rates. *Journal of Environmental Management* 79: 74–87. <https://doi.org/10.1016/j.jenvman.2005.05.017>
- Etter A, McAlpine C, Possingham H (2008) Historical patterns and drivers of landscape change in Colombia since 1500: A regionalized spatial approach. *Annals of the Association of American Geographers* 98: 2–23. <https://doi.org/10.1080/00045600701733911>
- Etter A, Wyngaarden WV (2000) Patterns of Landscape transformation in Colombia, with emphasis in the Andean region. *AMBIO: A Journal of the Human Environment* 29: 432–439. <https://doi.org/10.1579/0044-7447-29.7.432>
- Fariñas-Torres T, Pardiñas U, Chemisquy MA (2018) Los mamíferos de La Rioja, ocho décadas después de Yepes. *Revista del Museo Argentino de Ciencias Naturales nueva serie* 20: 123–135. <https://doi.org/10.22179/REVMACN.20.555>
- Ferreira MS, Vieira MV (2016) An efficient timing device to record activity patterns of small mammals in the field. *Mammalia* 80: 117–119. <https://doi.org/10.1515/mammalia-2014-0131>
- Fiedler E, Pavan SE, Paraense M, Author C, Tsuchiya MTN, Wilson DE, Percequillo AR, Jes P (2020) Museomics of tree squirrels: A dense taxon sampling of mitogenomes reveals hidden diversity, phenotypic convergence, and the need of a taxonomic overhaul. *BMC Evolutionary Biology* 20: 1–25. <https://doi.org/10.1186/s12862-020-01639-y>
- Garcés-Retrepo MF, Quintero-Angel A, Cuéllar N, Giraldo A (2016) Mammal diversity in an area with relicts of dry forest in the mid-Magdalena valley (Caldas, Colombia). *Revista de Ciencias* 20: 147–160. <https://doi.org/10.25100/rc.v20i2.4608>
- García-Herera L, Ramírez-Francel L, Reinoso-Flórez G (2019) Mamíferos del departamento del Tolima: distribución y estado de conservación. *Revista U.D.C.A Actualidad & Divulgación Científica* 22. <https://doi.org/10.31910/rudca.v22.n2.2019.1100>
- García-Herrera LV, Ramírez-Francel LA, Reinoso-Flórez G (2018) Potential distribution of *Sphaeronycteris toxophyllum* in Colombia and new record. *Therya* 9: 255–260. <https://doi.org/10.12933/therya-18-610>
- García FJ, Delgado-Jaramillo MI, Machado M, Aular L (2016) Mamíferos de la Sierra de Aroa, estado Yaracuy, Venezuela: listado taxonómico y la importancia de su conservación. *Memoria de la Fundación La Salle de Ciencias Naturales* 73: 17–34. [http://saber.ucv.ve/ojs/index.php/rev\\_mem/article/view/12298](http://saber.ucv.ve/ojs/index.php/rev_mem/article/view/12298)
- Gerkema MP, Davies WIL, Foster RG, Menaker M, Hut RA (2013) The nocturnal bottleneck and the evolution of activity patterns in mammals. *Proceedings of the Royal Society B: Biological Sciences* 208: 1–11. <https://doi.org/10.1098/rspb.2013.0508>
- Gerstner BE, Kass JM, Kays R, Helgen KM, Anderson RP (2018) Revised distributional estimates for the recently discovered olinguito (*Bassaricyon neblina*), with comments

- on natural and taxonomic history. *Journal of Mammalogy* 99: 321–332. <https://doi.org/10.1093/jmammal/gyy012>
- Haffer J (1967a) Speciation in Colombian forest birds west of the Andes. *American Museum Novitates* 2294: 1–57. <http://hdl.handle.net/2246/3087>
- Haffer J (1967b) Zoogeographical notes on the “nonforest” lowland bird faunas of north-western South America. *Hornero* 10: 315–333.
- Hammer Ø, Harper DAT, Ryan PD (2001) PAST: Paleontological statistics software package for education and data analysis. *Palaeontologia Electronica* 4: 1–9. [https://palaeo-electronica.org/2001\\_1/past/past.pdf](https://palaeo-electronica.org/2001_1/past/past.pdf)
- Hermelin M (2016) Landscapes and Landforms of Colombia. *World Geomorphological Landscapes*. Springer, Cham, 219 pp. <https://doi.org/10.1007/978-3-319-11800-0>
- Hernández-Camacho JH, Walschburger T, Quijano RO, Guerra AH (1992) Origen y distribución de la biota suramericana y colombiana. In: Halffter G (Ed.) *La Diversidad Biológica de Iberoamérica I*. Instituto de Ecología A.C., Xalapa, 55–104.
- Hernández-Canchola G, León-Paniagua L (2020) *Sturnira parvidens* (Chiroptera: Phyllostomidae). *Mammalian Species* 52(992): 57–70. <https://doi.org/10.1093/mspecies/seaa005>
- Hoffmann FG, Platt II RN, Mantilla-Meluk H, Medellín RA, Baker RJ (2019) Geographic and genetic variation in bats of the genus *Glossophaga*. In: Bradley RD, Genoways HH, Schmidly DJ, Bradley LC (Eds) *Special Publications Museum of Texas Tech University*, 71: 187–204. <https://www.depts.ttu.edu/nsrl/publications/downloads/SP71.pdf>
- Huck M, Juárez CP, Rotundo MA, Dávalos VM, Fernandez-Duque E (2017) Mammals and their activity patterns in a forest area in the Humid Chaco, northern Argentina. *Check List* 13(4): 363–378. <https://doi.org/10.15560/13.4.363>
- IDEAM (2001) Estudio ambiental de la cuenca Magdalena-Cauca y elementos para su ordenamiento territorial and Corporación Autónoma Regional del Río Grande de la Magdalena (CORMAGDALENA), Bogotá.
- Kattan GH, Franco P, Rojas V, Morales G (2004) Biological diversification in a complex region: A spatial analysis of faunistic diversity and biogeography of the Andes of Colombia. *Journal of Biogeography* 31: 1829–1839. <https://doi.org/10.1111/j.1365-2699.2004.01109.x>
- Larsen PA, Hooper SR, Bozeman MC, Pedersen SC, Genoways HH, Phillips CJ, Pumo DE, Baker RJ (2007) Phylogenetics and phylogeography of the *Artibeus jamaicensis* complex based on Cytochrome-*b* DNA sequences. *Journal of Mammalogy* 88: 712–727. <https://doi.org/10.1644/06-MAMM-A-125R.1>
- Lees AC, Naka LN, Aleixo A, Cohn-Haft M, Piacentini VDQ, Santos MPD, Silveira LF (2014) Conducting rigorous avian inventories: Amazonian case studies and a roadmap for improvement. *Revista Brasileira de Ornitologia* 22: 107–120. <https://doi.org/10.1007/BF03544240>
- Lim BK (2012) Biogeography of mammals from the Guianas of South America. In: Patterson BD, Costa LP (Eds) *Bones, clones, and biomes: the history and geography of recent neotropical mammals*. The University of Chicago Press, Chicago, 230–258. <https://doi.org/10.7208/chicago/9780226649214.003.0011>
- Lim BK, Loureiro LO, Garbino GST (2020) Cryptic diversity and range extension in the big-eyed bat genus *Chiroderma* (Chiroptera, Phyllostomidae). *ZooKeys* 918: 41–63. <https://doi.org/10.3897/zookeys.918.48786>

- Mantilla-Meluk H, Cadena GA, Jiménez-Ortega AM (2014) Historia de la mastozoología en Colombia: Pasado, presente y perspectivas. In: Ortega J, Martínez JL, Tirira DG (Eds) Historia de la mastozoología en Latinoamérica, las Guayanas y el Caribe. Editorial Murciélago Blanco y Asociación Ecuatoriana de Mastozoología, Quito y Mexico D.F., 153–174.
- Mena JL, Solari S, Carrera JP, Aguirre LF, Gómez H (2011) Small mammal diversity in the tropical Andes: an overview. In: Herzog SK, Martínez R, Jørgensen RM, Tiessen H (Eds) Climate change and biodiversity in the tropical Andes. Inter-American Institute for Global Change Research (IAI) and Scientific Committee on Problems of the Environment (SCOPE), São José dos Campos, 260–275.
- Minin ED, Slotow R, Hunter LTB, Pouzols FM, Toivonen T, Verburg PH, Leader-Williams N, Petracca L, Moilanen A (2016) Global priorities for national carnivore conservation under land use change. *Scientific Reports* 6: 1–9. <https://doi.org/10.1038/srep23814>
- Moreno-Bejarano LM, Álvarez-León R (2003) Fauna asociada a los manglares y otros humedales en el delta-estuario del río Magdalena, Colombia. *Revista de la Academia Colombiana de Ciencias Exactas, Físicas y Naturales* 27: 517–534. [http://www.accefyn.com/revista/Vol\\_27/105/4-FAUNA.pdf](http://www.accefyn.com/revista/Vol_27/105/4-FAUNA.pdf)
- Morrone JJ (2014) Biogeographical regionalisation of the neotropical region. *Zootaxa* 3782: 1–110. <https://doi.org/10.11646/zootaxa.3782.1.1>
- Mosquera-Guerra F, Trujillo F, Aya-Cuero C, Bolívar L, Valencia K, Vargas-Arboleda AF, Mantilla-Meluk H (2019) Mamíferos. In: Trujillo F, Anzola F (Eds) Biodiversidad en el departamento de Arauca. Gobernación de Arauca, Fundación Omacha y Fundación Ecollano, Bogotá, 233–273.
- Niemelä J (2000) Biodiversity monitoring for decision-making. *Annales Zoologici Fennici* 37: 307–317. <http://hdl.handle.net/1975/267>
- Paise G, Vieira EM, Prado PI (2020) Small mammals respond to extreme habitat fragmentation in the Brazilian Atlantic Forest according to the landscape continuum model. *Mammal Research* 65: 309–322. <https://doi.org/10.1007/s13364-019-00464-z>
- Poveda K, Sánchez-Palomino P (2004) Habitat use by the white-footed tamarin, *Saguinus leucopus*: a comparison between a forest-dwelling group and an urban group in Mariquita, Colombia. *Neotropical Primates* 12: 6–9.
- Pratas-Santiago LP, Gonçalves ALS, da Maia Soares AMV, Spironello WR (2016) The moon cycle effect on the activity patterns of ocelots and their prey. *Journal of Zoology* 299: 275–283. <https://doi.org/10.1111/jzo.12359>
- Racero-Casarrubia J, Ballesteros-Correa J, Pérez-Torres J (2015) Mamíferos del departamento de Córdoba-Colombia: historia y estado de conservación. *Biota Colombiana* 16: 128–148.
- Ramírez-Chaves HE, Pérez WA (2010) Mamíferos (Mammalia: Theria) del departamento del Cauca, Colombia. *Biota Colombiana* 11: 141–171. <http://revistas.humboldt.org.co/index.php/biota/article/view/237>
- Ramírez-Mejía AF, Sánchez F (2016) Activity patterns and habitat use of mammals in an Andean forest and a *Eucalyptus* reforestation in Colombia. *Hystrix, the Italian Journal of Mammalogy* 27: 1–7. <https://doi.org/10.4404/hystrix-27.2-11319>
- Rodríguez-Herrera B, Ramírez-Fernández JD, Villalobos-Chaves D, Sánchez R (2014) Actualización de la lista de especies de mamíferos vivientes de Costa Rica. *Mastozoología*



- Neotropical 21: 275–289. [http://www.scielo.org.ar/scielo.php?script=sci\\_arttext&pid=S0327-93832014000200008](http://www.scielo.org.ar/scielo.php?script=sci_arttext&pid=S0327-93832014000200008)
- Rojas-Díaz V, Reyes-Gutiérrez M, Alberico MS (2012) Mamíferos (Synapsida, Theria) del Valle del Cauca. Biota Colombiana 13: 99–116.
- Sánchez-Giraldo C, Daza JM (2017) Non-volant mammals from the protected areas associated to hydroelectric projects on the eastern slope of the northern Cordillera Central, Colombia. Check List 13: e2098. <https://doi.org/10.15560/13.2.2098>
- Mena DS, Böhm M, Arvanitidis C, Barber-Meyer S, Brummitt N, Chandler M, Chatzinikolaou E, Costello MJ, Ding H, García-Moreno J, Gill M, Haase P, Jones M, Juillard R, Magnusson WE, Martin CS, McGeoch M, Mihoub JB, Pettorelli N, Proença V, Peng C, Regan E, Schmiedel U, Simaika JP, Weatherdon L, Waterman C, Xu H, Belnap J (2017) Building capacity in biodiversity monitoring at the global scale. Biodiversity and Conservation 26: 2765–2790. <https://doi.org/10.1007/s10531-017-1388-7>
- Sikes RS, Gannon WL (2011) Guidelines of the American Society of Mammalogists for the use of wild mammals in research. Journal of Mammalogy 92: 235–253. <https://doi.org/10.1644/10-MAMM-F-355.1>
- Simmons NB, Cirranello AL (2020) Bat Species of the World: A taxonomic and geographic database. <https://batnames.org/>
- Sociedad Colombiana de Mastozoología (2020) Lista de referencia de especies de mamíferos de Colombia. Versión 1.2. Conjunto de datos/Lista de especies. <http://doi.org/10.15472/kl1whs>
- Solari S, Gómez-Ruiz D, Patiño-Castillo E, Villada-Cadavid T, López-M C (2020) Bat diversity of the Serranía de San Lucas (Bolívar and Antioquia), northern Colombia. Therya 11: 69–78. <https://doi.org/10.12933/therya-20-879>
- Solari S, Muñoz-Saba Y, Rodríguez-Mahecha JV, Defler TR, Ramírez-Chaves HE, Trujillo F (2013) Riqueza, endemismo y conservación de los mamíferos de Colombia. Mastozoología Neotropical 20: 301–365. <https://www.redalyc.org/pdf/457/45729294008.pdf>
- Torres DA, Rojas AE (2020) First record of Sanborn's bonneted bat, *Eumops hansae* (Molossidae), in Colombia. Mammalia 84(6): 595–599. <https://doi.org/10.1515/mammalia-2019-0140>
- Trujillo F, Mosquera-Guerra F, Díaz-Pulido A, Carvajal-Castro JD, Mantilla-Meluk H (2018) Mamíferos del Escudo Guayanés de Colombia. In: Lasso CA, Señaris JC (Eds) Fauna Silvestre del Escudo Guayanés (Colombia-Venezuela). Instituto de Investigación de Recursos Biológicos Alexander von Humboldt, Bogotá, 345–365.
- Velazco P, Patterson DB (2013) Diversification of the yellow-shouldered bats, genus *Sturnira* (Chiroptera, Phyllostomidae), in the new world tropics. Molecular Phylogenetics and Evolution 68: 683–698. <https://doi.org/10.1016/j.ympev.2013.04.016>
- Velazco PM, Patterson BD (2019) Small mammals of the Mayo River basin in northern Peru, with the description of a new species of *Sturnira* (Chiroptera: Phyllostomidae). Bulletin of the American Museum of Natural History 429: 1–70. <https://doi.org/10.1206/0003-0090.429.1.1>
- Zimbres B, Peres CA, Machado RB (2017) Terrestrial mammal responses to habitat structure and quality of remnant riparian forests in an Amazonian cattle-ranching landscape 206: 283–292. <https://doi.org/10.1016/j.biocon.2016.11.033>

## Supplementary material 1

### **Presence/absence matrix of 548 species of mammals from 28 regions in Central and South America**

Authors: Diego A. Torres, Abel Eduardo Rojas

Data type: Presence/absence matrix

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/neotropical.16.e57109.suppl1>